

Topological Incompleteness and Order Incompleteness of the Lambda Calculus

ANTONINO SALIBRA

Università Ca'Foscari di Venezia

A model of the untyped lambda calculus univocally induces a lambda theory (i.e., a congruence relation on λ -terms closed under α - and β -conversion) through the kernel congruence relation of the interpretation function. A semantics of lambda calculus is (*equationally*) *incomplete* if there exists a lambda theory which is not induced by any model in the semantics. In this paper we introduce a new technique to prove in a uniform way the incompleteness of all denotational semantics of lambda calculus which have been proposed so far, including the strongly stable one, whose incompleteness had been conjectured by Bastonero, Gouy and Berline. We apply this technique to prove the incompleteness of any semantics of lambda calculus given in terms of partially ordered models with a bottom element. This incompleteness removes the belief that partial orderings with a bottom element are intrinsic to models of the lambda calculus, and that the incompleteness of a semantics is only due to the richness of the structure of representable functions. Instead, the incompleteness is also due to the richness of the structure of lambda theories. Further results of the paper are: (i) an incompleteness theorem for partially ordered models with finitely many connected components (= minimal upward and downward closed sets); (ii) an incompleteness theorem for topological models whose topology satisfies a suitable property of connectedness; (iii) a completeness theorem for topological models whose topology is non-trivial and metrizable.

Categories and Subject Descriptors: F.4.1 [**Theory of Computation**]: Mathematical Logic—*Lambda calculus and related systems*; F.3.2 [**Theory of Computation**]: Semantics of Programming Languages—*Denotational semantics*

General Terms: Languages, Theory

Additional Key Words and Phrases: Lambda Calculus, Lambda Theories, Order and Topological Incompleteness, Orderability/Unorderability, Partially Ordered Models, Topological Models

1. INTRODUCTION

Lambda theories are equational extensions of the untyped lambda calculus that are closed under derivation. They arise by syntactical considerations, a lambda theory may correspond to a possible operational (observational) semantics of the lambda

Author's address: A. Salibra, Università Ca'Foscari di Venezia, Dipartimento di Informatica, Via Torino 155, 30172 Venezia, Italy salibra@dsi.unive.it.

Partially supported by MURST Cofin'01 COMETA Project.

This article is a revised and expanded version of two papers of the author which appeared respectively in the Proceedings of the Sixteenth Annual IEEE Symposium on Logic in Computer Science (LICS'01) (June 2001) [Salibra 2001a], and in the Proceedings of the Workshop on Böhm theorem: applications to Computer Science Theory (BOTH 2001) [Salibra 2001b].

Permission to make digital/hard copy of all or part of this material without fee for personal or classroom use provided that the copies are not made or distributed for profit or commercial advantage, the ACM copyright/server notice, the title of the publication, and its date appear, and notice is given that copying is by permission of the ACM, Inc. To copy otherwise, to republish, to post on servers, or to redistribute to lists requires prior specific permission and/or a fee.

© 2003 ACM 1529-3785/2003/0700-0001 \$5.00

calculus, as well as by semantic ones, a lambda theory may be induced by a model of lambda calculus through the kernel congruence relation of the interpretation function (see e.g. [Abramsky and Ong 1993; Barendregt 1984; Berline 2000]). Since the lattice of the lambda theories is a very rich and complex structure, syntactical techniques are usually difficult to use in the study of lambda theories. Therefore, semantic methods have been extensively investigated.

Computational motivations and intuitions justify Scott's view of models as partially ordered sets with a least element and of computable functions as monotonic functions over these sets. After Scott, mathematical models of the lambda calculus in various categories of domains (see e.g. [Abramsky 1991]) were classified into semantics according to the nature of their representable functions (see e.g. [Barendregt 1984; Berline 2000; Plotkin 1993]). Scott's continuous semantics [Scott 1972] is given in the category whose objects are complete partial orders and morphisms are Scott continuous functions. The stable semantics [Berry 1978] and the recent strongly stable semantics [Bucciarelli and Ehrhard 1991] are strengthenings of the continuous semantics. The stable semantics is given in the category of DI-domains with stable functions as morphisms, while the strongly stable one in the category of DI-domains with coherence, and strongly stable functions as morphisms. All these semantics are structurally and equationally rich in the sense that it is possible to build up 2^{\aleph_0} models in each of them inducing pairwise distinct lambda theories (see [Kerth 1998; Kerth 2001]). The problem of the equational richness is related to the problem of the equational completeness/incompleteness of a semantics: is the set of lambda theories induced by these semantics equal or strictly included within the set of all lambda theories?

The first incompleteness result was obtained by Honsell and Ronchi della Rocca [1992] for the continuous semantics. They proved, via a hard syntactical proof, that there exists a lambda theory which cannot be the theory of a continuous model. Following a similar method, Gouy [1995] proved the incompleteness of the stable semantics with a much harder syntactical proof. Semantic and more simple proofs of incompleteness for the continuous and stable semantics can be found in [Bastonerio and Gouy 1999]. Bastonerio [1998] provides an incompleteness result for the hypercoherence subsetting of the strongly stable semantics.

In this paper we introduce a new technique to prove in a uniform way the incompleteness of all the denotational semantics of the untyped lambda calculus which have been proposed so far. Most of the semantics which involve monotonicity with respect to some partial order fail to induce the lambda theory axiomatized by the equation $\Omega xx = \Omega$, where Ω is the usual λ -term which represents looping. The general technique used here for proving that a class \mathcal{C} of models is incomplete is the following:

- (1) Find a property P satisfied by all models in \mathcal{C} .
- (2) Find a lambda theory which is only induced by models not satisfying P .

We apply this technique to show the incompleteness of any lambda calculus semantics given in terms of partially ordered models with a bottom element. In particular we thus get the incompleteness of the (whole of) Bucciarelli-Ehrhard strongly stable semantics, which had been conjectured by Bastonerio and Gouy [1999] and by

Berline [2000], and, which was previously known only for the hypercoherence sub-setting [Bastoneo 1998].

The problem of the incompleteness of the semantics of lambda calculus is also related to the open problem of the order-incompleteness of lambda theories ([Selinger 1996]): does there exist a lambda theory which is not induced by any non-trivially partially ordered model? Such a problem can be also characterized in terms of connected components (= minimal upward and downward closed sets) of a partial ordering: a lambda theory \mathcal{T} is order-incomplete if, and only if, any partially ordered model inducing \mathcal{T} has an infinite number of connected components, each one containing *exactly* one element of the model. Towards an answer to the order-incompleteness problem, in this paper we show the existence of a lambda theory which is only induced by partially ordered models with an infinite number of connected components, each one containing *at most* one λ -term denotation.

Further results of the paper are two topological theorems. In the first one we show the incompleteness of any semantics given in terms of topological models whose topology satisfies a suitable property of connectedness. In the second one a proof is given of the completeness of the semantics of lambda calculus given in terms of topological models whose topology is non-trivial and metrizable.

Outline of the paper. The paper is organized as follows. In Section 2 we review the basic definitions of the lambda calculus and summarize definitions and results concerning topology that will be needed in the subsequent part of the paper. In particular, we recall the formal definition of a model of lambda calculus, and the topological notions of connectedness and separation.

A short and simple proof of incompleteness for the class of partially ordered models with a bottom element is presented in Section 3, while the order-incompleteness problem is discussed in Section 4, where we prove the incompleteness of the class of partially ordered models with finitely many connected components.

The property of closed-open-connectedness is introduced in Section 5, where a strong property of separation orthogonal to closed-open-connectedness is proven for suitable topological algebras. This result is used to obtain the topological incompleteness theorem.

In Section 6 we show that the semantics of lambda calculus given in terms of non-trivial metric spaces is complete.

Some open problems related to the incompleteness problem of the untyped lambda calculus are discussed in Section 7, while the last two sections of the paper are devoted to related work and conclusions.

2. PRELIMINARIES

To keep this article self-contained, we summarize some definitions and results that we need in the subsequent part of the paper. With regard to the lambda calculus we follow the notation and terminology of Barendregt [1984]. The main references for topological algebras are [Taylor 1977; Bentz 1999; Coleman 1996; Coleman 1997].

2.1 Topology

A topological space (A, τ) (we will occasionally avoid explicit mention of τ) is *non-trivial* if there are nonempty, proper subsets X and Y of A such that X is open

while Y is not. If A is a topological space then the closure of a subset U of A will be denoted by \bar{U} . Recall that $a \in \bar{U}$ if $U \cap V \neq \emptyset$ for every open neighborhood V of a . As a matter of notation, we write \bar{b} for the closure of set $\{b\}$.

For any space (A, τ) a preorder can be defined by

$$a \leq_{\tau} b \text{ iff } a \in \bar{b} \text{ iff } \forall U \in \tau (a \in U \Rightarrow b \in U).$$

In other words, $a \leq_{\tau} b$ if every neighbourhood of a is also a neighbourhood of b .

We have

$$\tau \text{ is } T_0 \text{ iff } \leq_{\tau} \text{ is a partial order.}$$

For any T_0 -space A the partial order \leq_{τ} is called *the specialization order* of τ . Note that any continuous map between T_0 -spaces is necessarily monotone and that the order is discrete (i.e., it satisfies $a \leq_{\tau} b$ iff $a = b$) iff A is a T_1 -space.

A space A is

- T_2 (or Hausdorff) if for all distinct $a, b \in A$ there exist open sets U and V with $a \in U$, $b \in V$ and $U \cap V = \emptyset$.
- $T_{2_{1/2}}$ (or completely Hausdorff) if for all distinct $a, b \in A$ there exist open sets U and V with $a \in U$, $b \in V$ and $\bar{U} \cap \bar{V} = \emptyset$.
- T_3 (or regular) if it is T_1 and, for every closed set $B \subseteq A$ and every $a \notin B$, there exist open sets U and V with $a \in U$, $B \subseteq V$ and $U \cap V = \emptyset$.

The previous axioms of separation can be relativized to pairs of elements. For example, a and b are $T_{2_{1/2}}$ -separable, if there exist open sets U and V with $a \in U$, $b \in V$ and $\bar{U} \cap \bar{V} = \emptyset$. T_2 -, T_1 -, T_0 -separability are similarly defined.

A *clopen* set is a subset of a space which is both open and closed. A space A is *connected* if A and \emptyset are the unique clopen sets of the space A . The connected component of an element a of a space A is the greatest connected subset of A including a . The connected components define a partition of the space A .

A *topological algebra* is a pair (\mathbf{A}, τ) where \mathbf{A} is an algebra and τ is a topology on the underlying set A with the property that each basic operation of \mathbf{A} is continuous with respect to τ .

2.2 Partial Orderings

Let (A, \leq) be a partially ordered set (poset). Two elements a and b of A are: (1) *comparable* if either $a \leq b$ or $b \leq a$; (2) *upward compatible* if they have an upper bound, i.e., there exists z such that $a \leq z$ and $b \leq z$; (3) *downward compatible* if they have a lower bound. A set $B \subseteq A$ is *upward direct* (*downward direct*) if, for all $a, b \in B$, a and b are upward (downward) compatible.

A set $B \subseteq A$ is an *upward* (*downward*) closed set if $b \in B$ and $b \leq a$ ($a \leq b$) imply $a \in B$. We use the notations $B \uparrow$ ($B \downarrow$, respectively) for the least upward (downward) closed set containing a subset B of A . We write $a \uparrow$ ($a \downarrow$, respectively) for $\{a\} \uparrow$ ($\{a\} \downarrow$).

The *inequality graph* of a poset (A, \leq) has the elements of A as nodes, while an edge connects two distinct nodes a and b if either $a < b$ or $b < a$. Two nodes are in the same connected component if they are either not distinct or joined by a path. The equivalence classes of the relation “to be in the same connected component”

define the partition of the inequality graph into *connected components*. A connected component can be also characterized as a minimal subset of A which is both upward closed and downward closed.

Given a poset (A, \leq) , we can find many T_0 -topologies τ on A for which \leq is the specialization ordering of τ (see [Johnstone 1982, Section II.1.8]). The maximal one with this property is the Alexandroff topology, which is constituted by the collection of all upward closed sets of A , i.e.,

$$U \text{ is an Alexandroff open iff } U = U\uparrow.$$

Then $a\uparrow$ is the least Alexandroff open set containing $a \in A$. A subset U of A is an Alexandroff closed set iff $U = U\downarrow$. A function is continuous w.r.t. the Alexandroff topology if, and only if, it is monotone. Every Alexandroff space is T_0 , but not T_1 , unless the order is the equality (if $x < y$, then every Alexandroff open set including x includes y too). Properties of the inequality graph of a poset can be expressed as topological properties of the Alexandroff topology. For example, the inequality graph of a poset is connected if, and only if, the Alexandroff topology is connected.

A *partially ordered algebra*, a *po-algebra* for short, is a pair (\mathbf{A}, \leq) where \mathbf{A} is an algebra and \leq is a partial order on A which makes the basic operations of \mathbf{A} monotone. Every po-algebra is a topological algebra w.r.t. the Alexandroff topology.

Every T_0 -topological algebra (\mathbf{A}, τ) is also a po-algebra relatively to the specialization order \leq_τ .

2.3 The Untyped Lambda Calculus

The two primitive notions of the lambda calculus are *application*, the operation of applying a function to an argument (expressed as juxtaposition of terms), and *lambda (functional) abstraction*, the process of forming a function from the “rule” that defines it. The set Λ of λ -terms of the lambda calculus over an infinite set X of variables is constructed as usual: every variable $x \in X$ is a λ -term; if M and N are λ -terms, then so are (MN) and $\lambda x.M$ for each variable x .

An occurrence of a variable x in a λ -term is *bound* if it lies within the scope of a lambda abstraction λx ; otherwise it is *free*. A λ -term without free variables is said to be *closed*. Λ^o denotes the set of closed λ -terms.

$M[x := N]$ is the result of substituting a λ -term N for all free occurrences of x in M subject to the usual provisos about renaming bound variables in M to avoid capture of free variables in N .

The axioms of the $\lambda\beta$ -calculus are as follows: M and N are arbitrary λ -terms and x, y are variables.

- (α) $\lambda x.M = \lambda y.M[x := y]$, for any variable y that does not occur free in M ;
- (β) $(\lambda x.M)N = M[x := N]$;

The rules for deriving equations from instances of (α) and (β) are the usual ones from equational calculus asserting that equality is a congruence for application and abstraction.

Lambda theories are equational extensions of the lambda calculus that are closed under derivation. If \mathcal{T} is a set of equations, then the formal system $\lambda + \mathcal{T}$ is obtained by adding to α - and β -conversion and to the equality rules the equations in \mathcal{T} as

new axioms. If \mathcal{T} is a set of equations, \mathcal{T}^+ is the set of equations provable in $\lambda + \mathcal{T}$. \mathcal{T} is a lambda theory if $\mathcal{T}^+ = \mathcal{T}$. As a matter of notation, $\mathcal{T} \vdash M = N$ stands for $\lambda + \mathcal{T} \vdash M = N$; this is also written as $M =_{\mathcal{T}} N$. A lambda theory \mathcal{T} is consistent if there exists at least an equation $M = N$ such that $\mathcal{T} \not\vdash M = N$.

$\lambda\beta$ is the minimal lambda theory. $\lambda\beta\eta$ is the lambda theory generated by the axiom scheme of η -conversion:

$$\lambda x.Mx = M \quad (x \text{ not free in } M).$$

$\lambda\beta\eta$ is the minimal extensional lambda theory, since it can be also generated by adding to α - and β -conversion and to the equality rules the following derivation rule of extensionality:

$$Mx = Nx \Rightarrow M = N \quad (x \text{ not free in } MN).$$

A closed λ -term M is *solvable* if, for every $P \in \Lambda$, there exist an integer n and $N_1, \dots, N_n \in \Lambda$ such that $MN_1 \dots N_n =_{\lambda\beta} P$. $M \in \Lambda$ is *solvable* if $\lambda x_1 \dots x_n.M$ is solvable, where x_1, \dots, x_n are all the variables occurring free in M . $M \in \Lambda$ is *unsolvable* if it is not solvable. Solvable λ -terms can be also characterized as follows: a λ -term M is solvable if, and only if, it has a *head normal form*, that is, $M =_{\lambda\beta} \lambda x_1 \dots x_n.yM_1 \dots M_k$ for some $n, k \geq 0$ and λ -terms M_1, \dots, M_k .

A lambda theory \mathcal{T} is called *semisensible* [Barendregt 1984, Def. 4.1.7(iii)] if $\mathcal{T} \not\vdash M = N$ whenever M is solvable and N is unsolvable. Every semisensible lambda theory is consistent.

Lambda theories may be defined according to syntactical considerations as well as to semantic ones. A lambda theory may be the equational theory of a model of the lambda calculus (see Section 2.5 below), or it may correspond to a possible observational semantics of the lambda calculus. In the second case, consider a subset \mathcal{Q} of Λ . If $M \in \Lambda$ then we write $M \in_{\lambda\beta} \mathcal{Q}$ if there exists P such that $M =_{\lambda\beta} P$ and $P \in \mathcal{Q}$. Define

$$M \approx_{\mathcal{Q}} N \text{ iff, for every context } C[\], C[M] \in_{\lambda\beta} \mathcal{Q} \Leftrightarrow C[N] \in_{\lambda\beta} \mathcal{Q}.$$

Then, the set $Th(\mathcal{Q})$, defined by

$$Th(\mathcal{Q}) = \{M = N : M, N \in \Lambda \text{ and } M \approx_{\mathcal{Q}} N\},$$

is a lambda theory, called the *contextual lambda theory* which arises from set \mathcal{Q} .

2.4 Combinatory Algebras

Combinatory logic is a formalism for writing expressions which denote functions. Combinators are designed to perform the same tasks as λ -terms, but without using bound variables. Schönfinkel and Curry discovered that a formal system of combinators, having the same expressive power of the lambda calculus, can be based on only two primitive combinators.

We begin with the definition of a basic notion in combinatory logic and lambda calculus. An algebra $\mathbf{C} = (C, \cdot, \mathbf{k}, \mathbf{s})$, where \cdot is a binary operation and \mathbf{k}, \mathbf{s} are constants, is called a *combinatory algebra* (see [Curry and Feys 1958]) if it satisfies the following identities (as usual the symbol \cdot is omitted, and association is to the left):

$$\mathbf{k}xy = x; \quad \mathbf{s}xyz = xz(yz).$$

\mathbf{k} and \mathbf{s} are called the *basic combinators*. In the equational language of combinatory algebras the derived combinators \mathbf{i} and $\mathbf{1}$ are defined as follows: $\mathbf{i} := \mathbf{s}\mathbf{k}\mathbf{k}$ and $\mathbf{1} := \mathbf{s}(\mathbf{k}\mathbf{i})$. Hence, every combinatory algebra satisfies the identities $\mathbf{i}x = x$ and $\mathbf{1}xy = xy$.

A combinatory algebra \mathbf{C} is *degenerate* if C is a singleton set.

A function $f : C \rightarrow C$ is *representable* in the combinatory algebra \mathbf{C} if there exists an element $c \in C$ such that $cz = f(z)$ for all $z \in C$. If this last condition is satisfied, we say that c represents f in \mathbf{C} .

Two elements $c, d \in C$ are called *extensionally equal* if they represent the same function in \mathbf{C} . For example, the elements c and $\mathbf{1}c$ are extensionally equal for every $c \in C$. The combinator $\mathbf{1}$ will be used in the next Subsection to select a canonical representative inside the class of all elements d extensionally equal to a given element $c \in C$.

2.5 Lambda Models

Although lambda calculus has been the subject of research by logicians since the early 1930's, its model theory developed only much later, following the pioneering model construction made by Dana Scott. The notion of an *environment model* (the name is due to Meyer [1982]) is described by Meyer as “the natural, most general formulation of what might be meant by mathematical models of the untyped lambda calculus”. The drawback of environment models is that they are higher-order structures. However, there exists an intrinsic characterization (up to isomorphism) of environment models as an elementary class of combinatory algebras called *λ -models* [Barendregt 1984, Def. 5.2.7]. They were first axiomatized by Meyer [1982] and independently by Scott [1980]; the axiomatization, while elegant, is not equational.

We now define the notion of a λ -model. Let \mathbf{C} be a combinatory algebra and let \bar{c} be a new symbol for each $c \in C$. Extend the language of the lambda calculus by adjoining \bar{c} as a new constant symbol for each $c \in C$. Let $\Lambda^o(C)$ be the set of closed λ -terms with constants from C . The interpretation of terms in $\Lambda^o(C)$ with elements of C can be defined by induction as follows (for all $M, N \in \Lambda^o(C)$ and $c \in C$):

$$|\bar{c}|_{\mathbf{C}} = c; |(MN)|_{\mathbf{C}} = |M|_{\mathbf{C}}|N|_{\mathbf{C}}; |\lambda x.M|_{\mathbf{C}} = \mathbf{1}m,$$

where $m \in C$ is any element representing the following function $f : C \rightarrow C$:

$$f(c) = |M[x := \bar{c}]|_{\mathbf{C}}, \quad \text{for all } c \in C. \quad (1)$$

The drawback of the previous definition is that, if \mathbf{C} is an arbitrary combinatory algebra, it may happen that the function f is not representable. The axioms of an elementary subclass of combinatory algebras, called *λ -models* or models of the lambda calculus, were expressly chosen to make coherent the previous definition of interpretation (see [Meyer 1982], [Scott 1980], [Barendregt 1984, Def. 5.2.7]).

The *Meyer-Scott axiom* is the most important axiom in the definition of a λ -model. In the first-order language of combinatory algebras it takes the following form

$$\forall x \forall y (\forall z (xz = yz) \Rightarrow \mathbf{1}x = \mathbf{1}y)$$

and it makes the combinator $\mathbf{1}$ an inner choice operator. Indeed, given any c , the element $\mathbf{1}c$ is in the same equivalence class as c w.r.t. extensional equality; and, by Meyer-Scott axiom, $\mathbf{1}c = \mathbf{1}d$ for every d extensionally equal to c . Thus, the set $Y = \{c : cz = f(z) \text{ for all } z \in C\}$ of elements representing the function f defined in (1) admits $\mathbf{1}m$ as a canonical representative and this does not depend on the choice of $m \in Y$.

As a matter of notation, we write $\mathbf{C} \models M = N$ for $|M|_{\mathbf{C}} = |N|_{\mathbf{C}}$.

For every λ -model \mathbf{C} , the lambda theory generated by the set $\{M = N : M, N \in \Lambda^o, \mathbf{C} \models M = N\}$ is called *the equational theory of \mathbf{C}* and is denoted by $Th(\mathbf{C})$. We say that a lambda theory \mathcal{T} is *induced* by a λ -model \mathbf{C} (or that \mathbf{C} *induces* \mathcal{T}) if $\mathcal{T} = Th(\mathbf{C})$.

The *term model* $\mathcal{M}_{\mathcal{T}}$ of a lambda theory \mathcal{T} (see [Barendregt 1984, Def. 5.2.11]) consists of the set of the equivalence classes of λ -terms modulo the lambda theory \mathcal{T} together with the operation of application on the equivalence classes. By Cor. 5.2.13(ii) in [Barendregt 1984] $\mathcal{M}_{\mathcal{T}}$ is a λ -model which induces the lambda theory \mathcal{T} .

A *partially ordered model*, a po-model for short, is a pair (\mathbf{C}, \leq) , where \mathbf{C} is a λ -model and \leq is a partial order on C which makes the application operator of \mathbf{C} monotone. In particular a po-model is a po-algebra relatively to the language of combinatory algebras. A po-model (\mathbf{C}, \leq) is *non-trivial* if the partial order \leq is not discrete (i.e., $a < b$ for some $a, b \in C$).

A *topological model* of the lambda calculus is a topological algebra (\mathbf{C}, τ) , where \mathbf{C} is a λ -model.

3. PARTIALLY ORDERED MODELS WITH A BOTTOM ELEMENT

As already mentioned, there are computational motivations for taking ordered structures with a bottom element as models of the lambda calculus. The question now arises whether every lambda theory is induced by a po-model with a bottom element. In this Section we negatively answer to this question: we exhibit a lambda theory that does not arise from such a kind of models.

Consider the lambda theory Π axiomatized by

$$\Omega xx = \Omega,$$

where $\Omega \equiv (\lambda x.xx)(\lambda x.xx)$. Π is semisensible (Ω and Ωxx are both unsolvable) and then consistent.

The following Lemma will be repeatedly used in the remaining part of this article.

LEMMA 3.1. *Let M and N be arbitrary λ -terms. Then we have*

$$\Pi \vdash M = N \Leftrightarrow \Pi \vdash \Omega MN = \Omega.$$

PROOF. Let \rightarrow_{Π} be the compatible closure of the binary relation R on Λ defined as follows:

$$R = \{(\Omega MN, \Omega) : M, N \in \Lambda, \Pi \vdash M = N\}.$$

Let $\rightarrow_{\beta\Pi}$ be the reduction rule $\rightarrow_{\beta} \cup \rightarrow_{\Pi}$. Recall that \rightarrow_{β}^* is the reflexive, transitive closure of β -reduction \rightarrow_{β} .

We divide the proof into two Claims.

Claim 3.2. The relation $\rightarrow_{\beta\Pi}$ is Church-Rosser.

It is sufficient to verify that (a) the reflexive, transitive closure \rightarrow_{Π}^* of \rightarrow_{Π} satisfies the diamond property (see [Barendregt 1984, Lemma 3.2.2]); (b) the relations \rightarrow_{β}^* and \rightarrow_{Π}^* commute (see [Barendregt 1984, Def. 3.3.4]). The conclusion follows from the Hindley-Rosen Lemma (see [Barendregt 1984, Prop. 3.3.5]).

Claim 3.3. Π is the lambda theory generated by conversion $\cong_{\beta\Pi}$ from $\rightarrow_{\beta\Pi}$, i.e.,

$$\Pi \vdash M = N \text{ iff } M \cong_{\beta\Pi} N.$$

If $\Omega MN \rightarrow_{\Pi} \Omega$ then we have that $\Pi \vdash M = N$ by definition of \rightarrow_{Π} , so that $\Pi \vdash \Omega MN = \Omega$ follows from $\Pi \vdash \Omega xx = \Omega$. Then it is obvious that $M \cong_{\beta\Pi} N$ implies $\Pi \vdash M = N$. For the opposite direction, it is sufficient to consider that $\Omega xx \rightarrow_{\Pi} \Omega$ for the unique axiom $\Omega xx = \Omega$ of Π .

We now conclude the proof of the lemma. If $\Pi \vdash \Omega MN = \Omega$ then $\Omega MN \cong_{\beta\Pi} \Omega$, so that by Claim 3.2 there is a reduction $\Omega MN \rightarrow_{\beta\Pi}^* \Omega$. This is possible only if ΩMN is a Π -redex, i.e., if $\Pi \vdash M = N$. \square

We formally define the concepts of equational completeness/incompleteness of a class of λ -models.

Definition 3.4. A class \mathcal{Q} of λ -models is called (*equationally*) *complete* if every lambda theory is induced by some λ -model of \mathcal{Q} . If this is not the case, \mathcal{Q} is called *incomplete*.

The author is indebted to Gordon Plotkin for providing the short and simple proof of Thm. 3.5 below. The original proof of the author was based on topological methods (see [Salibra 2001a], [Salibra 2001b]).

THEOREM 3.5. *Any semantics of lambda calculus given in terms of po-models with a bottom element is incomplete.*

PROOF. Assume, by the way of contradiction, that the lambda theory Π is induced by a po-model (\mathbf{C}, \leq) with a bottom element \perp . From $\Pi \vdash \Omega xx = \Omega$ it follows that:

$$\mathbf{C} \models \Omega xx = \Omega. \tag{2}$$

Let $\mathbf{k} \equiv \lambda xy.x$ and $\mathbf{i} \equiv \lambda y.y$ (it is indeed sufficient that \mathbf{k} and \mathbf{i} are two distinct $\beta\eta$ -normal forms). By monotonicity and by (2) the following conditions hold in \mathbf{C} :

$$\Omega \mathbf{k} \mathbf{i} \geq \Omega \perp \perp = \Omega,$$

and then

$$\Omega = \Omega \Omega \Omega \leq \Omega(\Omega \mathbf{k} \mathbf{i})\Omega \leq \Omega(\Omega \mathbf{k} \mathbf{i})(\Omega \mathbf{k} \mathbf{i}) = \Omega.$$

So, we conclude:

$$\Omega(\Omega \mathbf{k} \mathbf{i})\Omega = \Omega.$$

Then by applying twice Lemma 3.1 we get first

$$\Omega \mathbf{k} \mathbf{i} = \Omega,$$

and second

$$\mathbf{k} = \mathbf{i}.$$

This contradicts the consistency of the lambda theory Π . \square

The following corollary solves the open problem of the incompleteness of the strongly stable semantics.

COROLLARY 3.6. *The continuous, stable, and strongly stable semantics of lambda calculus are incomplete.*

We say that a semantics *omits* a lambda theory \mathcal{T} if \mathcal{T} is not induced by any model in the semantics. In [Salibra 2001a] the author has shown that continuous, stable and strongly stable semantics omit a continuum of lambda theories. In fact, by applying a well-known theorem by Visser (see [Visser 1980], [Barendregt 1984, Thm. 17.1.10]), it is possible to get a continuum of distinct lambda theories satisfying the conditions:

$$\Omega xx = \Omega; \quad \Omega(\Omega \mathbf{k} \mathbf{i})\Omega \neq \Omega.$$

From the proof of Thm. 3.5 it follows that any semantics given in terms of po-models with a bottom element omits each of these lambda theories.

4. ORDER INCOMPLETENESS

In the previous Section we have given a negative answer to the question whether every lambda theory arises as the theory of a po-model with a bottom element. One may now relax this question by asking: does every lambda theory arise as the equational theory of a non-trivial po-model? The answer to this difficult question is still unknown. Selinger [1996] gave a syntactical characterization, in terms of the so-called generalized Mal'cev operators, of the order-incomplete lambda theories (i.e., the theories not induced by any non-trivial po-model). A lambda theory \mathcal{T} is order-incomplete if, and only if, there exist a natural number $n \geq 1$ and a sequence M_1, \dots, M_n of closed λ -terms such that the following Mal'cev conditions are satisfied:

$$x =_{\mathcal{T}} M_1 x y y; \quad M_i x x y =_{\mathcal{T}} M_{i+1} x y y; \quad M_n x x y =_{\mathcal{T}} y \quad (1 \leq i < n).$$

Plotkin and Simpson have shown that the above Mal'cev conditions are inconsistent with lambda calculus for $n = 1$, while Plotkin and Selinger obtained the same result for $n = 2$ (see [Selinger 1997]). It is an open problem whether n can be greater than or equal to 3.

The problem of order-incompleteness can be also characterized as follows: a lambda theory \mathcal{T} is order-incomplete if, and only if, the inequality graph (see Section 2.2) of any po-model inducing \mathcal{T} has an infinite number of connected components, each one containing exactly one element of the model. In this Section we give a partial solution to the problem of order-incompleteness by showing the following result: the inequality graph of any po-model inducing the lambda theory Π (defined in Section 3) has an infinite number of connected components, each one containing at most one λ -term denotation. This result implies the incompleteness of any semantics of lambda calculus given in terms of po-models with a finite number of connected components.

We begin the technical part of this Section by introducing the notion of a semisubtractive algebra. These algebras satisfy a very weak form of subtractivity. We recall that the notion of subtractivity in Universal Algebra was introduced by Ursini [1994]: an algebra is *subtractive* if it satisfies the identities

$$s(x, x) = 0; \quad s(x, 0) = x \quad (3)$$

for some binary term s and constant 0 . Subtractive algebras abound in classical algebras and in algebraic logic since term s simulates part of subtraction. If we interpret the binary operator “ s ” as subtraction “ $-$ ” and we use the infix notation, then we can rewrite the above identities (3) as $x - x = 0$ and $x - 0 = x$.

Definition 4.1. An algebra \mathbf{A} is *semisubtractive* if there exist a binary term $s(x, y)$ and a constant 0 in the similarity type of \mathbf{A} such that

$$s(x, x) = 0.$$

Example 4.2. Let \mathbf{C} be a λ -model inducing the lambda theory Π defined in Section 3. Define

$$0 \equiv \Omega; \quad s(x, y) \equiv \Omega xy.$$

From $\Pi \vdash \Omega xx = \Omega$ it easily follows that $\mathbf{C} \models s(x, x) = 0$. Then \mathbf{C} is a semisubtractive algebra.

A semisubtractive algebra is *trivial* if $s(x, y) = 0$ for all x and y . Every semisubtractive po-algebra, which has both a bottom element \perp and a top element \top , is trivial:

$$0 = s(\perp, \perp) \leq s(x, y) \leq s(\top, \top) = 0.$$

In the following we always assume that a semisubtractive algebra is non-trivial.

As a matter of notation, ω denotes the first infinite ordinal, i.e., the set of natural numbers.

Definition 4.3. Let \mathbf{A} be a semisubtractive algebra. The *subtraction sequence* $(c_n)_{n \in \omega}$ of a pair $(a, b) \in A^2$ is defined by induction as follows:

$$c_1 = s(a, b); \quad c_{n+1} = s(c_n, 0). \quad (4)$$

We say that (a, b) has *order* $k \in \omega$ if $c_k \neq 0$, while (a, b) has *order* ω if $c_k \neq 0$ for all $k \in \omega$.

Notice that, if a pair (a, b) has order k , then $a \neq b$ and (a, b) has order n for all $n \leq k$. In fact, if $c_n = 0$, then $c_{n+1} = s(c_n, 0) = s(0, 0) = 0$, that is, if some $c_n = 0$, then all c_j with $j > n$ are 0.

If we interpret the binary operator “ s ” as subtraction “ $-$ ” and we use the infix notation, then we can rewrite the elements of the sequence (c_n) as follows:

$$c_1 = a - b; \quad c_2 = (a - b) - 0; \quad c_3 = ((a - b) - 0) - 0; \quad \text{and so on.}$$

In the ring of integers all elements of the sequence are equal and they are all different from 0 if a and b are distinct.

Recall from Section 2.2 the definition of the inequality graph of a po-algebra.

LEMMA 4.4. *If (\mathbf{A}, \leq) is a semisubtractive po-algebra and a and b are joined by an edge in the inequality graph of A , then $s(a, b) = 0$.*

PROOF.

$$a \leq b \Rightarrow 0 = s(a, a) \leq s(a, b) \leq s(b, b) = 0,$$

and similarly for $b \leq a$. \square

LEMMA 4.5. *Let (\mathbf{A}, \leq) be a semisubtractive po-algebra. If $a, b \in A$ are joined by a path of length k (in the inequality graph of A), then the pair (a, b) cannot have order k .*

PROOF. The proof is by induction on the length k of the path. If $k = 1$ the conclusion follows from Lemma 4.4.

Assume now a and b to be connected by a path of length $k + 1$. Then there exists an element d such that a and d are connected by a path of length k , and d, b are joined by an edge (without loss of generality, we may assume $d < b$). Let (c_n) be the subtraction sequence of (a, b) and (r_n) be the subtraction sequence of (a, d) . By the induction hypothesis the pair (a, d) does not have order k , so that

$$r_k = 0.$$

Since $d < b$, by an easy induction we have that $r_n \leq c_n$ for all n . Then we get

$$0 = r_k \leq c_k,$$

that is, 0 and c_k are equal or joined by an edge in the inequality graph of \mathbf{A} . The conclusion $c_{k+1} \equiv s(c_k, 0) = 0$ follows from Lemma 4.4. \square

COROLLARY 4.6. *Let (\mathbf{A}, \leq) be a semisubtractive po-algebra. If a pair $(a, b) \in A^2$ has order ω , then a and b are in distinct connected components.*

In the next theorem we characterize the po-models inducing the lambda theory Π defined in Section 3. We start with a Lemma.

LEMMA 4.7. *Let M and N be arbitrary λ -terms. Consider the sequence of λ -terms defined by induction as follows:*

$$c_1 \equiv \Omega MN; \quad c_{n+1} \equiv \Omega(c_n)\Omega$$

(c_i will become a subtraction sequence later). Then we have

$$\Pi \not\vdash M = N \Rightarrow \Pi \not\vdash c_n = \Omega \text{ for all } n.$$

PROOF. Trivial by induction on n , using Lemma 3.1. \square

THEOREM 4.8. *Let (\mathbf{C}, \leq) be a po-model of lambda calculus. If $Th(\mathbf{C}) = \Pi$ then the inequality graph of \mathbf{C} has an infinite number of connected components. Moreover, if $\Pi \not\vdash M = N$ then the interpretations of M and N in \mathbf{C} are in distinct connected components.*

PROOF. By Example 4.2 \mathbf{C} is a semisubtractive algebra, so that by Lemma 4.7 any pair (M, N) of two non- Π -equivalent closed λ -terms has order ω . Then by Cor. 4.6 M and N are in distinct connected components of the inequality graph of \mathbf{C} . Since Π is consistent there are infinitely many pairwise non- Π -equivalent

closed λ -terms (for example, the non- $\beta\eta$ -equivalent normal forms), and then infinite distinct connected components. \square

THEOREM 4.9. (Order Incompleteness Theorem) *Any semantics of the untyped lambda calculus, given in terms of po-models which have only finitely many connected components, is incomplete.*

PROOF. By Thm. 4.8. \square

COROLLARY 4.10. *Any semantics of lambda calculus given in terms of po-models which are semilattices, lattices, complete partial orderings, or which have a top or a bottom element, is incomplete.*

5. TOPOLOGICAL INCOMPLETENESS

In this Section we show the incompleteness of any semantics of lambda calculus, given in terms of topological models satisfying a suitable property of connectedness called *closed-open-connectedness*. The proof is based on a general theorem of separation for topological algebras.

5.1 A Topological Theorem of Separation

We recall that separation axioms in topology stipulate the degree to which distinct points may be separated by open sets or by closed neighbourhoods of open sets. In Thm. 5.2 below we prove that in every semisubtractive T_0 -topological algebra every pair of elements of order 3 is $T_{2_{1/2}}$ -separable. We were inspired with Bentz and Coleman for the idea of this theorem and for the techniques used in its proof (see [Bentz 1999; Coleman 1996; Coleman 1997]).

THEOREM 5.1. *If (\mathbf{A}, τ) is a semisubtractive T_0 -topological algebra, then every pair $(a, b) \in A^2$ of order 2 is T_2 -separable.*

PROOF. Let (c_n) be the subtraction sequence of (a, b) . Since \mathbf{A} is a po-algebra with respect to the specialization ordering \leq_τ of τ (see Section 2.2), then from the hypothesis $c_2 = s(c_1, 0) \neq 0$ and from Lemma 4.4 it follows that c_1 and 0 are incomparable w.r.t. \leq_τ , i.e., neither $c_1 \leq_\tau 0$ nor $0 \leq_\tau c_1$. Then there exists an open neighbourhood U of c_1 such that $0 \notin U$.

From $s(a, b) = c_1 \in U$ and from the continuity of s , it follows that there exist two open neighbourhoods V, W , of a and b respectively, such that $s(V, W) \subseteq U$. If there is an element $e \in V \cap W$, then $0 = s(e, e) \in U$. This contradicts the hypothesis on U . Hence we have $V \cap W = \emptyset$. \square

THEOREM 5.2. *If (\mathbf{A}, τ) is a semisubtractive T_0 -topological algebra, then every pair $(a, b) \in A^2$ of order 3 is $T_{2_{1/2}}$ -separable.*

PROOF. Let (c_n) be the subtraction sequence of (a, b) and (d_n) be the subtraction sequence of $(c_1, 0)$. It is obvious that we have $d_n = c_{n+1}$ for all n and, hence, that $(c_1, 0)$ has order 2. By applying Thm. 5.1 we get an open neighbourhood V' of c_1 and an open neighbourhood W' of 0 such that $V' \cap W' = \emptyset$.

Since s is continuous and $s(a, b) = c_1 \in V'$, there exist two other open sets V and W containing a and b , respectively, such that $s(V, W) \subseteq V'$. The sets V and W will be the right sets for the conclusion of the theorem. Since s is continuous the

pre-image of $\overline{V'}$ under the map s is closed. From $s(V, W) \subseteq V' \subseteq \overline{V'}$ the pre-image of $\overline{V'}$, which is closed, contains $V \times W$, so $s(\overline{V}, \overline{W}) \subseteq \overline{V'}$.

We now prove that $\overline{V} \cap \overline{W} = \emptyset$. Assume, by the way of contradiction, that there is $e \in \overline{V} \cap \overline{W}$. Since $s(\overline{V}, \overline{W}) \subseteq \overline{V'}$ it follows that $0 = s(e, e) \in \overline{V'}$. But by definition of closure of a set this is possible only if, for every open neighbourhood Z of 0 , we have that $Z \cap V' \neq \emptyset$. But this contradicts our initial choice of V' and W' as two open neighbourhoods of c_1 and 0 respectively with empty intersection. \square

5.2 Closed-Open-Connectedness

Connectedness axioms in topology examine the structure of a topological space in an orthogonal way with respect to separation axioms. They deny the existence of certain subsets of a topological space with properties of separation. In this Section we introduce a strong property of connectedness, called closed-open-connectedness, which is orthogonal to the property of $T_{2_{1/2}}$ -separability, and it is satisfied by a topological space if there exist no $T_{2_{1/2}}$ -separable elements.

Definition 5.3. We say that a space is *closed-open-connected*, co-connected for short, if it has no disjoint closures of open sets. In other words, if, for all open sets U and V , we have that $\overline{U} \cap \overline{V} \neq \emptyset$.

We recall that a space with no disjoint open sets is called *hyperconnected*, while a space with no disjoint closed sets is called *ultraconnected* (see [Steen and Seebach 1978, Section 4]). Then we have the following implications:

$$\text{hyperconnectedness} \Rightarrow \text{co-connectedness} \Rightarrow \text{connectedness}$$

and

$$\text{ultraconnectedness} \Rightarrow \text{co-connectedness} \Rightarrow \text{connectedness}.$$

Then closed-open-connectedness is a sort of meeting point between ultraconnectedness and hyperconnectedness.

The following proposition provides a wide class of topological spaces whose topology is co-connected.

PROPOSITION 5.4. *Let (X, τ) be a T_0 -topological space, whose specialization order \leq_τ satisfies the following property: every pair of nodes of the inequality graph of (X, \leq_τ) is joined by a path of length less or equal to 3. Then (X, τ) is co-connected.*

PROOF. Assume, by the way of contradiction, that there exist elements $a, b \in X$ and open sets U and V such that $a \in U$, $b \in V$ and $\overline{U} \cap \overline{V} = \emptyset$. Then a and b are incomparable w.r.t. the specialization ordering, otherwise either $a \in V$ or $b \in U$. By hypothesis a and b are joined by a path of length less or equal to 3. We have four possibilities.

- (1) $(\exists c) a < c > b$. In this case $c \in U \cap V$.
- (2) $(\exists c) a > c < b$. Then we have $c \in \overline{a} \subseteq \overline{U}$ and $c \in \overline{b} \subseteq \overline{V}$.
- (3) $(\exists c, d) a < c > d < b$. Then $d \in \overline{c} \subseteq \overline{U}$ and $d \in \overline{b} \subseteq \overline{V}$.
- (4) $(\exists c, d) a > c < d > b$. Then $c \in \overline{d} \subseteq \overline{V}$ and $c \in \overline{a} \subseteq \overline{U}$.

All these possibilities contradict our assumption. \square

COROLLARY 5.5. *Every T_0 -topological space (X, τ) , whose specialization order either admits a bottom (top) element or makes X an upward (downward) direct set, is co-connected.*

In particular complete partial orderings (including continuous, stable and strongly stable models) with the Scott topology are co-connected.

We cannot improve Prop. 5.4. The following counter-example provides a space, whose topology is not co-connected, and such that every pair of nodes of the inequality graph associated with the specialization order \leq_τ is joined by a path of length less or equal to 4. Let $X = \{a, b, c, d, e\}$ be a set partially ordered as follows: $a > b < c > d < e$. Then X is a topological space w.r.t. the Alexandroff topology (see Section 2.2). X is not co-connected, since $a\uparrow\downarrow = \{a, b\}$ and $e\uparrow\downarrow = \{d, e\}$ are disjoint closures of the open sets $a\uparrow$ and $e\uparrow$.

5.3 The Topological Incompleteness Theorem

As a consequence of the theorem of separation of Section 5.1, we show that the topological models of the lambda theory Π are not co-connected. The topological incompleteness theorem follows from this result.

Recall from Section 2.5 that a topological model of the lambda calculus is a topological algebra (\mathbf{C}, τ) , where \mathbf{C} is a λ -model.

THEOREM 5.6. *Let (\mathbf{C}, τ) be a T_0 -topological model of the lambda calculus. If $Th(\mathbf{C}) = \Pi$ then τ is not co-connected.*

PROOF. We recall from Example 4.2 that \mathbf{C} is a semisubtractive algebra. Let $\mathbf{k} \equiv \lambda xy.x$ and $\mathbf{i} \equiv \lambda y.y$ (it is indeed sufficient that \mathbf{k} and \mathbf{i} are two distinct $\beta\eta$ -normal forms). Then the subtraction sequence of (\mathbf{k}, \mathbf{i}) has order ω by Lemma 4.7, so that \mathbf{k} and \mathbf{i} are $T_{2_{1/2}}$ -separable by Thm. 5.2. \square

We conclude this Section with the topological incompleteness theorem.

THEOREM 5.7. (Topological Incompleteness Theorem) *Any semantics of the lambda calculus given in terms of co-connected T_0 -topological models is incomplete.*

PROOF. From Thm. 5.6. \square

We now analyse the relationships between the order incompleteness theorem and the topological incompleteness theorem.

Recall from Section 2.1 that a topological model, whose topology is T_0 but not T_1 , has a non-trivial specialization order ($a < b$ for some a, b). Then from Thm. 4.9 and from Thm. 5.7 it follows the incompleteness of any semantics of lambda calculus given in terms of T_0 -topological models, whose topology is either co-connected or admits a specialization order with a finite number of connected components.

Partial orderings can be viewed as topological spaces w.r.t. the Alexandroff topology (see Section 2.2). Then the order incompleteness theorem can be also expressed as follows (see [Salibra 2001b]): any semantics of lambda calculus given in terms of Alexandroff topological models with a finite number of connected components is incomplete.

All semantics we have shown incomplete in Cor. 4.10 are constituted by po-models whose Alexandroff topology is co-connected (see Cor. 5.5). Then the incompleteness

of continuous, stable and strongly stable semantics also follows from the topological incompleteness theorem.

The following natural questions arise:

- (Q1) Is there a po-model of lambda calculus whose Alexandroff topology is connected but not co-connected?
- (Q2) Is there a po-model of lambda calculus whose Alexandroff topology has a finite number (greater than 1) of connected components?
- (Q3) Is there a non-trivial co-connected T_1 -topological model of lambda calculus? In other words, is there a co-connected topological model of lambda calculus, whose specialization order is the equality relation?

6. COMPLETENESS

In this Section we positively answer the following question: is there a class of λ -models with a “reasonable” topology which is a complete semantics of lambda calculus? In the main result of this Section we show that the semantics of lambda calculus, given in terms of topological models whose topology is non-trivial and metrizable, is complete. We conclude the section by discussing some negative results about the existence of topological models in categories of topological spaces.

If (A, d) is a metric space, then τ_d is the topology on A defined by d . A *metrizable model* of the untyped lambda calculus is a topological model whose topology is non-trivial and metrizable.

THEOREM 6.1. (Metric Completeness Theorem) *The semantics of the untyped lambda calculus given in terms of metrizable models is complete.*

PROOF. Let \mathcal{T} be an arbitrary consistent lambda theory. The term model $\mathcal{M}_{\mathcal{T}}$ of \mathcal{T} defined in Section 2.5 is a λ -model [Barendregt 1984, Cor. 5.2.13(ii)] such that $Th(\mathcal{M}_{\mathcal{T}}) = \mathcal{T}$. We will show that it is possible to define on $\mathcal{M}_{\mathcal{T}}$ a metric d in such a way that $(\mathcal{M}_{\mathcal{T}}, \tau_d)$ is a metrizable model. Consider the closed term model $\mathcal{M}_{\mathcal{T}}^c$ of \mathcal{T} , that is the subalgebra of $\mathcal{M}_{\mathcal{T}}$ generated by the equivalence classes of closed λ -terms. Notice that $\mathcal{M}_{\mathcal{T}}^c$ is not in general a λ -model (see [Barendregt 1984, Cor. 5.2.13(i)]).

Let X be the set of variables of lambda calculus. Then $\mathcal{M}_{\mathcal{T}}$ is the free extension of $\mathcal{M}_{\mathcal{T}}^c$ by X in the variety of combinatory algebras (see e.g. [Pigozzi and Salibra 1998, Section 3.3], [Salibra 2000, Section 4.2]), that is, $\mathcal{M}_{\mathcal{T}}$ satisfies the following universal mapping property: for every homomorphism $h : \mathcal{M}_{\mathcal{T}}^c \rightarrow \mathbf{A}$ from $\mathcal{M}_{\mathcal{T}}^c$ into a combinatory algebra \mathbf{A} and every mapping $g : X \rightarrow A$ there exists a unique homomorphism $f : \mathcal{M}_{\mathcal{T}} \rightarrow \mathbf{A}$ of combinatory algebras extending both h and g .

Since $\mathcal{M}_{\mathcal{T}}$ is a free algebra, then we can apply the following construction due to Świerczkowski [1964] (see also [Taylor 1977, Thm. 2.1] and [Coleman 1997, Section 3]). Let (X, d_0) be an arbitrary metric space whose universe X is the set of variables of lambda calculus. By applying Świerczkowski’s theorem, it is possible to extend the metric d_0 to a metric d on the free algebra $\mathcal{M}_{\mathcal{T}}$ in such a way that the application operator is continuous w.r.t. τ_d and $d(x, y) = d_0(x, y)$ for all $x, y \in X$. The conclusion of the theorem follows because $(\mathcal{M}_{\mathcal{T}}, \tau_d)$ is a metrizable model of \mathcal{T} . \square

Are there other classes of topological models which are complete semantics of the lambda calculus? In order to answer this question, it is necessary to consider

the properties of separation of the topological models of the lambda theory Π (see Thm. 5.6). We recall that many authors tried to find lambda calculus models in Cartesian closed categories of topological spaces. Abramsky (see [Mislove 1998, Thm. 5.11]) and Plotkin (see [Mislove 1998, Thm. 5.14]) have shown respectively that there exists no nondegenerate model of the lambda calculus in the category of posets and monotone mappings, and in the category of complete ultrametric spaces and non-expansive mappings. Hofmann and Mislove [1995] have shown that the category of k -spaces and continuous maps has no nondegenerate, compact T_2 -topological model. A k -space is a topological space in which a subset is open if and only if its intersection with each compact subset of the space is open in the subspace. The following problem is still open.

(Q4) (Hoffman-Mislove) Is there a lambda calculus model in the category of k -spaces?

Notice that every topological λ -model, in which all the continuous selfmaps of the model are representable (as in the category of k -spaces), must have a connected topology because of the existence of fixed points. If a topology is not connected, then there exists a clopen (open and closed) set U and two elements a and b such that $a \in U$ and $b \notin U$. Then the function f , defined by

$$f(x) = b \text{ if } x \in U; \quad f(x) = a \text{ if } x \notin U,$$

is continuous without fixed points. The following natural question now arises:

(Q5) Is the semantics of lambda calculus, given in terms of connected topological models, complete?

7. ORDERABILITY/UNORDERABILITY

In this section we present some further results and discuss some open problems related to the incompleteness problem of the untyped lambda calculus.

If every model of lambda calculus were nontrivially orderable, then the order-incompleteness problem (discussed in Section 4) would admit a negative answer. However, Selinger [1996] has recently shown that the term models $\mathcal{M}_{\lambda\beta}$ and $\mathcal{M}_{\lambda\beta\eta}$ of the lambda theories $\lambda\beta$ and $\lambda\beta\eta$ are unorderable (i.e., they do not admit a nontrivial compatible partial order). As a consequence of the theorem of separation in Section 5.1, we now show that the term model \mathcal{M}_{Π} of the lambda theory Π is also unorderable.

Definition 7.1. An algebra \mathbf{A} is *strongly semisubtractive* if it is semisubtractive and satisfies the following condition:

$$s(x, y) = 0 \Rightarrow x = y. \quad (5)$$

PROPOSITION 7.2. *A strongly semisubtractive T_0 -topological algebra \mathbf{A} is unorderable.*

PROOF. Assume, by the way of contradiction, that there exists a non-trivial compatible partial order on \mathbf{A} . Let $a, b \in A$ be distinct elements with $a < b$. Then from Lemma 4.4 it follows that $s(a, b) = 0$, which implies $a = b$ by (5). Contradiction. \square

COROLLARY 7.3. \mathcal{M}_Π is strongly semisubtractive, and hence unorderable.

PROOF. By Lemma 4.7 and by Prop. 7.2. \square

PROPOSITION 7.4. A strongly semisubtractive T_0 -topological algebra \mathbf{A} is $T_{2_{1/2}}$.

PROOF. From (5) it follows that every pair of distinct elements of A has order ω . Then the conclusion follows from Thm. 5.2. \square

It is not possible to improve the conclusion of Prop. 7.4, as shown by the following example suggested to the author by E. Jabara.

Example 7.5. Let R be the set of real numbers, $A = \{(x, y) \mid x, y \in R, y \geq 0\}$ be the upper halfplane of R^2 and τ_0 be the usual Euclidean topology on A . For every $(a, 0) \in A$ and for every real number $\epsilon > 0$ define

$$K_a(\epsilon) = \{(x, y) \in A \mid (x - a)^2 + y^2 < \epsilon, y > 0\} \cup \{(a, 0)\}.$$

If τ is the least topology on A containing all the elements of τ_0 and all $K_a(\epsilon)$ (for every $a, \epsilon \in R$ with $\epsilon > 0$), then it is possible to verify that (A, τ) is a $T_{2_{1/2}}$ -space but not a T_3 -space. The binary operation \star on A , defined by

$$(x_1, y_1) \star (x_2, y_2) = (x_1 - x_2, |y_1 - y_2| + 1),$$

is continuous w.r.t. the topology τ , and it simulates part of the subtraction (with “0” represented by the pair $(0, 1)$):

$$(x_1, y_1) \star (x_2, y_2) = (0, 1) \text{ iff } x_1 = x_2, y_1 = y_2.$$

Then (A, \star, τ) is a strongly semisubtractive topological algebra whose topology is not T_3 .

The classification of the λ -models into orderable/unorderable models can be refined as follows. For every model \mathbf{C} , let $T_i^{\mathbf{C}}$ ($i = 0, 1, 2, 2_{1/2}$) be the set of all topologies τ on C which make (\mathbf{C}, τ) a T_i -topological model. It is obvious that in general we have

$$T_0^{\mathbf{C}} \supseteq T_1^{\mathbf{C}} \supseteq T_2^{\mathbf{C}} \supseteq T_{2_{1/2}}^{\mathbf{C}}.$$

We recall from Section 2.1 that a topology with a non-trivial specialization order (we have $a < b$ for some a, b) would be T_0 yet not T_1 , so that

$$\mathbf{C} \text{ is unorderable iff } T_0^{\mathbf{C}} = T_1^{\mathbf{C}}.$$

We say that a lambda theory \mathcal{T} is of (topological) type i ($i = 0, 1, 2, 2_{1/2}$) if the term model of \mathcal{T} satisfies $T_0^{\mathcal{M}\mathcal{T}} = T_i^{\mathcal{M}\mathcal{T}}$. All lambda theories are of type 0; the lambda theory \mathcal{B} , generated by equating two lambda terms if they have the same Böhm tree, is not of type 1 (see [Barendregt 1984]). $\lambda\beta$ and $\lambda\beta\eta$ are of type 1 by Selinger’s result, while Π is of type $2_{1/2}$ by Cor. 7.3 and by Prop. 7.4. Then some natural questions arise:

(Q6) Is $\lambda\beta$ ($\lambda\beta\eta$) of type 2?

(Q7) Is $\lambda\beta$ ($\lambda\beta\eta$) of type $2_{1/2}$?

Selinger has shown in [Selinger 1997; Selinger 1996] that the problem of the order-incompleteness is also related to the following question by Plotkin [1996]:

(Q8) Is there an absolutely unorderable combinatory algebra?

A combinatory algebra is absolutely unorderable if it cannot be embedded in any non-trivially partially ordered combinatory algebra. In the last result of this Section we give a partial positive answer to Plotkin's question (see Lusin and Salibra [200X] for a different approach to this problem).

We recall that a quasi-variety is a class of algebras axiomatized by quasi-identities (i.e., equational implications with a finite number of equational premises).

Definition 7.6. Let \mathcal{V} be a quasi-variety of algebras and let $\mathbf{C} \in \mathcal{V}$. We say that \mathbf{C} is *absolutely unorderable in \mathcal{V}* if, for every embedding $\mathbf{C} \hookrightarrow \mathbf{D} \in \mathcal{V}$, \mathbf{D} is unorderable.

PROPOSITION 7.7. *Let \mathcal{SD} be the quasi-variety of combinatory algebras axiomatized by $\Omega xy = \Omega \Leftrightarrow x = y$. Then the term model \mathcal{M}_Π of the lambda theory Π is absolutely unorderable in \mathcal{SD} .*

PROOF. \mathcal{M}_Π belongs to \mathcal{SD} because of Cor 7.3. Then the conclusion follows from Prop. 7.2 since all algebras in \mathcal{SD} are strongly semisubtractive. \square

8. RELATED WORK

In this Section we review the classic results of incompleteness for the continuous and stable semantics. The first incompleteness result was obtained by Honsell and Ronchi della Rocca [1992] for the continuous semantics. They extended to all continuous models arising from an inverse limit construction an approximation theorem introduced by Hyland and Wadsworth to study the Scott model D_∞ (see [Barendregt 1984]). The general approximation theorem was applied to prove that the lambda theory induced by the Park continuous model \mathbf{P} is included within the contextual lambda theory $Th(\Lambda^\circ)$ induced by the set Λ° of closed λ -terms (see Section 2.3 for the definition of contextual lambda theory). Then the authors prove the incompleteness of the continuous semantics by showing that, whenever the theory of a continuous model is included in $Th(\Lambda^\circ)$, then it is *strictly* included in $Th(\Lambda^\circ)$. Namely, there exist two λ -terms X and Z such that

$$Th(\Lambda^\circ) \vdash X = Z, \tag{6}$$

while

$$Th(\mathbf{C}) \not\vdash X = Z \tag{7}$$

for every continuous model \mathbf{C} such that $Th(\mathbf{C}) \subseteq Th(\Lambda^\circ)$. The complexity of the proof comes from the fact that (6) is established syntactically, while (7) is a consequence of the existence of non-sequential functions within the set of all Scott continuous functions.

Following a method similar to that of Honsell and Ronchi della Rocca [1992], Gouy [1995] proved the incompleteness of the stable semantics. Other more semantic proofs of incompleteness for the continuous, stable and hypercoherence semantics (that is a subclass of the strongly stable semantics introduced by Ehrhard [1993]) can be found in [Bastoneo 1998; Bastoneo and Gouy 1999] and are briefly described in the following. The Park model \mathbf{P} was first defined in the framework of continuous semantics. It is a variant of the Scott model D_∞ , but with a very

different equational theory. This model has a stable analogue \mathbf{P}_s (which was defined by Honsell and Ronchi della Rocca [1990]), and a strongly stable analogue \mathbf{P}_{fs} (defined by Bastonero and Gouy [1999]). It is possible to give a semantic proof of incompleteness for the continuous semantics by using the Park stable model \mathbf{P}_s . Bastonero and Gouy [1999] prove that the λ -terms X and Z , considered by Honsell and Ronchi della Rocca [1992], are distinct in all continuous models \mathbf{C} satisfying the following conditions:

- (i) \mathbf{C} is extensional,
- (ii) $\Omega\Omega = \Omega$,
- (iii) $\lambda x.\Omega(\Omega x) = \Omega$,
- (iv) $\Omega \neq \lambda x.\Omega$.

The Park stable model \mathbf{P}_s satisfies the above conditions (i)-(iv) and equalizes X and Z (see [Bastonero and Gouy 1999, Thm 4.1]). This proves that no continuous model has the theory of \mathbf{P}_s .

The incompleteness of the stable semantics was proven by Bastonero and Gouy in a similar way. They show that the same λ -terms X and Z are distinct in all stable models \mathbf{C} satisfying the following conditions:

- (a) \mathbf{C} is extensional,
- (b) $\Omega\Omega = \Omega$,
- (c) $\Omega(\lambda x.\Omega) \neq \Omega$.

The Park strongly stable model \mathbf{P}_{fs} satisfies the above three conditions (a)-(c) and equalizes X and Z (see [Bastonero and Gouy 1999, Thm 5.1]). This proves that no stable model has the theory of \mathbf{P}_{fs} .

The same technique was applied by Bastonero [1998] to prove the incompleteness of the hypercoherence semantics. He shows that there exist two λ -terms F and G which are distinct in all hypercoherent models satisfying a suitable set \mathcal{P} of constraints. Then, by using the technique of forcing, Bastonero builds a continuous model \mathbf{C} which satisfies the constraints \mathcal{P} and equalizes F and G . This proves that no hypercoherent model has the theory of \mathbf{C} .

9. CONCLUSION

In this paper we have introduced a new technique to prove in a uniform way the incompleteness of all the denotational semantics of lambda calculus which have been proposed so far, including the strongly stable one, whose incompleteness had been conjectured by Bastonero and Gouy [1999] and by Berline [2000]. We apply this technique to prove the incompleteness of any semantics of lambda calculus given in terms of the following classes of models: (i) Partially ordered models with a bottom element; (ii) Partially ordered models with a finite number of connected components; (iii) Co-connected topological models. Our incompleteness theorems remove the belief that complete partial orders are intrinsic to models of the lambda calculus, and that the incompleteness of the continuous semantics is only due to the richness of the structure of the Scott continuous functions. Instead, the incompleteness is also due to the richness of the structure of the lambda theories.

ACKNOWLEDGMENTS

The author wishes to thank Gordon Plotkin and the referees for helpful comments and suggestions.

REFERENCES

- ABRAMSKY, S. 1991. Domain theory in logical form. *Annals of Pure and Applied Logic* 51, 1–77.
- ABRAMSKY, S. AND ONG, C. 1993. Full abstraction in the lazy λ -calculus. *Information and Computation* 105, 159–267.
- BARENDREGT, H. 1984. *The lambda calculus: Its syntax and semantics*. Number 103 in Studies in Logic. North-Holland Publishing Co., Amsterdam.
- BASTONERO, O. 1998. Equational incompleteness and incomparability results for λ -calculus semantics. Manuscript.
- BASTONERO, O. AND GOUY, X. 1999. Strong stability and the incompleteness of stable models of λ -calculus. *Annals of Pure and Applied Logic* 100, 247–277.
- BENTZ, W. 1999. Topological implications in varieties. *Algebra Universalis* 42, 9–16.
- BERLINE, C. 2000. From computation to foundations via functions and application: The λ -calculus and its webbed models. *Theoretical Computer Science* 249, 81–161.
- BERRY, G. 1978. Stable models of typed lambda-calculi. In *Proc. 5th Int. Coll. on Automata, Languages and Programming*. LNCS, vol. 62. Springer-Verlag, Berlin.
- BUCCIARELLI, A. AND EHRHARD, T. 1991. Sequentiality and strong stability. In *Proceedings of the Sixth Annual IEEE Symposium on Logic in Computer Science*. IEEE Computer Society Publications, Los Alamitos, CA, USA, 138–145.
- COLEMAN, J. 1996. Separation in topological algebras. *Algebra Universalis* 35, 72–84.
- COLEMAN, J. 1997. Topological equivalents to n -permutability. *Algebra Universalis* 38, 200–209.
- CURRY, H. AND FEYS, R. 1958. *Combinatory Logic, Vol. I*. North-Holland Publishing Co., Amsterdam.
- EHRHARD, T. 1993. Hypercoherences: a strongly stable model of linear logic. *Mathematical Structures in Computer Science* 2, 365–385.
- GOUY, X. 1995. Etude des théories équationnelles et des propriétés algébriques des modèles stables du λ -calcul. Ph.D. thesis, Université de Paris 7, Paris, France.
- HOFMANN, K. AND MISLOVE, M. 1995. All compact hausdorff lambda models are degenerate. *Fundamenta Informaticae* 22, 23–52.
- HONSELL, F. AND RONCHI DELLA ROCCA, S. 1990. Reasoning about interpretation in qualitative λ -models. In *Programming Concepts and Methods*, M. Broy and C. Jones, Eds. Elsevier Science Publishing Co., Amsterdam, 505–521.
- HONSELL, F. AND RONCHI DELLA ROCCA, S. 1992. An approximation theorem for topological λ -models and the topological incompleteness of λ -calculus. *Journal Computer and System Science* 45, 49–75.
- JOHNSTONE, P. 1982. *Stone Spaces*. Cambridge University Press, Cambridge.
- KERTH, R. 1998. Isomorphism and equational equivalence of continuous lambda models. *Studia Logica* 61, 403–415.
- KERTH, R. 2001. On the construction of stable models of λ -calculus. *Theoretical Computer Science* 269, 23–46.
- LUSIN, S. AND SALIBRA, A. 200X. A note on absolutely unorderable combinatory algebras. *Journal of Logic and Computation*. to appear.
- MEYER, A. 1982. What is a model of the lambda calculus? *Information and Control* 52, 87–122.
- MISLOVE, M. 1998. Topology, domain theory and theoretical computer science. *Topology and its Applications* 89, 3–59.
- PIGOZZI, D. AND SALIBRA, A. 1998. Lambda abstraction algebras: coordinatizing models of lambda calculus. *Fundamenta Informaticae* 33, 149–200.
- PLOTKIN, G. 1993. Set-theoretical and other elementary models of the λ -calculus. *Theoretical Computer Science* 121, 351–409.

- PLOTKIN, G. 1996. On a question of H. Friedman. *Information and Computation* 126, 74–77.
- SALIBRA, A. 2000. On the algebraic models of lambda calculus. *Theoretical Computer Science* 249, 197–240.
- SALIBRA, A. 2001a. A continuum of theories of lambda calculus without semantics. In *Proceedings of the Sixteenth Annual IEEE Symposium on Logic in Computer Science (LICS'01)*. IEEE Computer Society Publications, Los Alamitos, CA, USA, 334–343.
- SALIBRA, A. 2001b. Towards lambda calculus order-incompleteness. In *Workshop on Böhm theorem: applications to Computer Science Theory (BOTH 2001)*. Electronics Notes in Theoretical Computer Science, vol. 50. Elsevier Science Publishing Company, Amsterdam, 147–160.
- SCOTT, D. 1972. Continuous lattices. In *Toposes, Algebraic geometry and Logic*, F. Lawvere, Ed. Number 274 in LNCS. Springer-Verlag, Berlin, 97–136.
- SCOTT, D. 1980. Lambda calculus: some models, some philosophy. In *The Kleene Symposium*, H. K. J. Barwise and K. Kunen, Eds. Number 101 in Studies in Logic. North-Holland Publishing Co., Amsterdam, 97–136.
- SELINGER, P. 1996. Order-incompleteness and finite lambda models. In *Proceedings of the Eleventh Annual IEEE Symposium on Logic in Computer Science*. IEEE Computer Society Publications, Los Alamitos, CA, USA.
- SELINGER, P. 1997. Functionality, polymorphism, and concurrency: a mathematical investigation of programming paradigms. Ph.D. thesis, University of Pennsylvania.
- STEEN, L. AND SEEBACH, J. 1978. *Counterexamples in topology*. Springer-Verlag, Berlin.
- ŚWIERCZKOWSKI, S. 1964. Topologies in free algebras. *Proc. London Math. Soc.* 3, 566–576.
- TAYLOR, W. 1977. Varieties obeying homotopy laws. *Canad. Journal Math.* 29, 498–527.
- URSINI, A. 1994. On subtractive varieties, I. *Algebra Universalis* 31, 204–222.
- VISSER, A. 1980. Numerations, λ -calculus and arithmetic. In *To H.B. Curry: Essays on Combinatory Logic, Lambda-Calculus and Formalism*, J. Hindley and J. Seldin, Eds. Academic Press, New York, 259–284.

Received October 2001; revised August 2002; accepted October 2002